NAG5-713 IN-47-CR 8356

UNIVERSITY OF CALIFORNIA Los Angeles

North Atlantic Weather Regimes: A Synoptic Study of Phase Space

A thesis submitted in partial satisfaction of the requirements for the degree Master of Science in Atmospheric Sciences

by

Anna Karin Orrhede

1990

(NASA-CR-188162) NORTH ATLANTIC WEATHER REGIMES: A SYNOPTIC STUDY OF PHASE SPACE M.S. Thesis (California Univ.) 8 pCSCL 04B N91-21673

Unclas G3/47 0008356

The thesis of Anna Karin Orrhede is approved

Akio Arakawa

David Neelin

Michael Ghil, Committee Chair

University of California, Los Angeles
1990

TABLE OF CONTENTS

	Acknowledgements v
	Abstractvi
1.	Introduction
	1.1 Dynamical background
	1.2 Combining statistics and dynamics 5
	1.3 A new approach
2.	Data
3 .	The trajectory minimization method
	3.1 EOFs and the phase space of weather 15
	3.2 Depicting the trajectory
	3.3 Finding weather regimes
	3.4 The regimes over the North Atlantic and northwestern Europe
	3.5 The classification of events
4.	Persistent events and the behaviour of the trajectory 28
	4.1 The path in phase space
	4.2 Duration of the events
	4.3 Regime transitions

	4.4 The trajectory within and between events 39
5 .	Synoptic study of the regimes
	5.1 The blocking regime 46
	a. Classification of the activities 50
	b. Actions and reactions 53
	c. Action – reaction sequences 55
	d. The oscillation and balance
	e. Trajectory behaviour 60
	5.2 The zonal regime
	a. Event activities
	b. Interactions
	c. Trajectory behaviour
	5.3 The Greenland anticyclone regime 72
	a. Event activities
	b. Responses and interactions
	c. Trajectory behaviour
	5.4 The Atlantic ridge regime 80
	a. Important factors in the flow 81
	b. Interactions in different situations 83
	c. Trajectory behaviour
6.	Synoptic study of the transitions

	6.1 The BL→GA transition	• •	86
	6.2 The ZO→BL transition	•• !	90
	6.3 The ZO→AR transition	!	94
7.	Conclusions	9	98
	7.1 The study of phase space	9	99
	7.2 The case studies	. 10)1
	7.3 Extension and future work	10)4
	References	. 10)6
	Table captions	11	0
	Tables	11	.1
	Figure captions	11	3
	Figures	19	1

ACKNOWLEDGEMENTS

I wish to thank Professor Michael Ghil for advising and inspiring me in this work. Also Dr. Robert Vautard has been a great help in introducing me to this subject and providing me with data as well as material. I am most greatful to Dr. Cécile Penland's support, personally and professionally, through discussions and meetings. She also supplied me with some additional data crucial for this work. Thanks are also addressed to Professors Akio Arakawa and David Neelin for contributing with their extensive knowledge, and to Dr. Kingtse Mo for interesting discussions. Johan has stood by me in every imaginable way and has given me lots of useful advice concerning general physics and computing. This work has been supported partly by NASA Grant NAG5-713 and by NSF Grant ATM86-15424.

ABSTRACT OF THE THESIS

North Atlantic Weather Regimes: A Synoptic Study of Phase Space

by

Anna Karin Orrhede

Master of Science in Atmospheric Sciences

University of California, Los Angeles, 1990

Professor Michael Ghil, Chair

In the phase space of weather, low-frequency variability (LFV) of the atmosphere can be captured in a large-scale subspace, where a trajectory connects consecutive large-scale weather maps, thus revealing flow changes and recurrence. Using this approach, Vautard (1989) applied the trajectory-speed minimization method (Vautard and Legras, 1988) to atmospheric data. From 37 winters of 700 mb geopotential height anomalies over the North Atlantic and the adjacent land masses, four persistent and recurrent weather patterns, interpreted as weather regimes, were discernable: a blocking regime, a zonal regime, a Greenland anticyclone regime and an Atlantic ridge regime.

In this study, we investigate these regimes further in terms of

maintenance and transitions. A regime survey unveils preferences regarding event durations and precursors for the onset or break of an event. The transition frequencies between regimes vary, and together with the transition times, suggest the existence of easier transition routes. These matters are more systematically studied using complete synoptic map sequences from a number of events. In the maintenance of an event, oscillations between different balancing forces are observed, in connection with recurring nonlinear interaction sequences. Also the transition routes seem to contain particular evolution sequences leading to the next quasi-stationary flow pattern.